

# **YOUTH SUICIDE AWARENESS & PREVENTION**

**Guidance and Information  
for Teachers and Lecturers**

# YOUTH SUICIDE AWARENESS & PREVENTION

## Guidance and Information for Teachers and Lecturers

---

This document forms part of the Youth Suicide Prevention Ireland Four Steps to Help Programme Training Pack. This document is copyrighted to the authors and licensed to Youth Suicide Prevention Ireland Publications Limited on an exclusive basis. The Four Steps to Help Programme is republished by kind permission of Alan McKelvie and Anthony Philpott. **Four Steps to Help Programme Copyright © Alan McKelvie and Anthony Philpott 2020 – 2024. All rights reserved. E&OE**

This publication may not be copied, reproduced or changed in any way without the express written consent of the publishers. You must obtain written permission before editing or changing this document. Selling without prior written consent is prohibited. In all cases this notice must remain intact. This work cannot be retailed or resold without the publisher's written consent. This work is registered with the UK Copyright Service as Copyright Witness and the rights of the Author are protected under International Copyright Law. UKCS Registration Number 324197. ([www.copyrightservice.co.uk](http://www.copyrightservice.co.uk)). Published by Youth Suicide Prevention Ireland Publications Limited.

**Extracts from "Youth Self-Harm and Suicide" republished by kind permission of Doctor Keith Holmes. Copyright © Doctor Keith Holmes 2020 – 2024. All Rights reserved.**

**Some materials used in this manual are republished by kind permission of the Samaritans of Rhode Island, USA. ([www.samaritansri.org](http://www.samaritansri.org)) Those materials have been adapted to the Irish Curriculum and localised into Irish English. Copyright © Samaritans of Rhode Island 2020 – 2024. All Rights reserved.**

All information herein is provided as an information resource about suicide prevention and research activities only. It is not the intention of Youth Suicide Prevention Ireland Publications Limited to provide specific treatment or care advice. Youth Suicide Prevention Ireland Publications Limited does not provide certification or accreditation for any counselling organisation, individual practitioners or treatment or care models that users may find out about through our information or publications or other materials provided by or through us. Youth Suicide Prevention Ireland Publications Limited does not endorse or recommend any of the service providers, agencies or organisations listed on any leaflet or publication provided by us or any other organisation. In no event shall liability exist to any user (or any third party who obtains access to the information) under any law for any loss or damage (including but not limited to direct, indirect, economic and consequential damage) in respect of any matter whatsoever related to any information in any website or leaflet or publication nor for the use of the information nor for actions taken or not taken as a result of information contained in this or any other websites leaflets or publications (including, without limitation, loss or damage caused by negligence). The material contained in this leaflet is provided for general information purposes only and does not constitute care, treatment or other professional advice. Due to the general nature of the information provided in the leaflet, should you require specific help, you should seek the assistance of an appropriately trained professional person. While every care has been taken in the preparation of the information contained in this leaflet, due to the nature of the information and the processes of publication, the information may not be complete, correct or up to date. Information contained in this leaflet which has been provided by recognised third parties is provided in good faith and no liability shall exist for any errors or omissions contained.

# CONTENTS

<b>Introduction</b> .....	4
The role of the teacher.....	4
The role of education.....	5
<b>An introduction to youth self-harm and suicide</b> .....	6
The Four Steps to Help Programme.....	6
Why awareness and vigilance are so important.....	7
Fighting stigma.....	8
<b>Frequently Asked Questions</b> .....	8
<b>Myths about suicide</b> .....	10
<b>What should I be watching for?</b> .....	13
Some of the underlying causes of youth suicide and self-harm.....	13
Cyber Bullying.....	14
Suicide Clusters.....	15
<b>Warning signs and behavioural changes</b> .....	16
Main Indicators.....	17
General Warning Signs.....	18
Teen specific warning signs.....	18
General risk factors.....	19
Teen specific risk factors.....	21
The Dangerous Calm.....	21
<b>Basic guidelines on how to help</b> .....	22
Tolerant and accepting.....	22
Respect.....	22
Not judging or trivialising.....	22
Allow the young person to talk.....	23
Don't guarantee confidentiality.....	23
Listening.....	23
Discussion does not cause suicide.....	23
Under 18s.....	23
Getting support.....	23
Helping students – Do's.....	24
Helping Students – Don'ts.....	24
Mental Health Awareness Programmes.....	24
Availability of staff.....	24
Providing students with coping skills.....	25
What can I do for my students?.....	25
<b>The role of the listener</b> .....	26
General guidance.....	26
Active Listening.....	27
Characteristics of good active listening.....	27
Attitude of the listener.....	28
<b>If you cannot help a student</b> .....	29
<b>Policy matters</b> .....	29
Student Protection.....	29
Self-protection.....	29
<b>Types of intervention Resources</b> .....	30
<b>Summary</b> .....	31

# INTRODUCTION

Teachers enjoy a unique role in the moulding and development of their students. It is unique in one sense—the teacher may never know the impact that he or she has had on a given student. Because it is difficult to ascertain the degree of influence, the teacher may falsely conclude that there is no influence, or very little, outside of intellectual skills; or our PE teachers might conclude that they have little influence outside of the development of motor skills. Whether you wish to admit it or not, we are constantly imparting affective skills.

Paradoxically, it is in the last area that teachers may feel insecure or may even deny having any influence. Our students need us in more ways than we can ever know. We need to teach them how to think, but we must also teach them how to deal with emotions. We need to teach them the accumulated wisdom of centuries, but we must also teach them to cope with the here and now. We must teach them about historical and literary characters, but we must also teach them to deal with the people in their environment. We must teach them the skills necessary to earn a living, but we must also teach them about death and dying. In order to teach them, however, we must learn ourselves.

We can no longer perceive our roles only as dispensers of knowledge. The students sitting in front of us are not mere receptacles for ideas, facts, values, trends, data, or events. Those sitting in front of us are caught up in the pressure-cooker of modern life. Students of all ages are sometimes seething with feelings that are incomprehensible, with a sense of helplessness, hopelessness, and no sense of the future. Setting a course in uncharted territory, their feelings may be emotionally debilitating, with ideas that confuse rather than clarify, and with demands that would defy Hercules. Just as adults can be overwhelmed so can students, and classroom activities may be the last thing they are thinking about.



## The role of the teacher

The modern adolescent, bereft of any apparent solution to his problems and seemingly cut off from sharing his pain with another human being, may break under the pressure and opt for the most permanent solution - death. And who but teachers are in the best position to observe students individually or in groups, to sense the emotions seething under the surface, to monitor subtle changes in behaviour, and to realise that the student in reality might be a hurting human being?

But if the teacher lacks knowledge about suicide - its causes and its warning signs - the teacher very well could be unaware that a life and death drama might be developing. It is too late to rewrite the lesson plans of life once the student rips up the original copy.

This is not meant to reduce teachers to a state of paranoia about suicide. The intent is to call the teachers' attention to the need for self-analysis about their own attitudes toward suicide and its victims, and to the need for an awareness of the signals that students may be sending out about their inner turmoil.

Chad Varan, founder of The Samaritans, put all our minds at ease when he argued that one doesn't need a degree or diploma to help another human being. He said, "What is needed to save another human being from death ... is a compassionate heart, the willingness to accept, to pay attention, to care." Who among our teaching staffs does not have these characteristics? We teach, so we care. All we need is knowledge that will direct our caring in the right direction.



In talking to teachers informally, we have learned that many of us are willing and ready to befriend our students. However, we are sometimes frustrated by a system that does not allow us the time and the freedom to sit and listen to our students who are crying out for help, acceptance, and love.

The following guide deals with death by suicide. Certainly not a very pleasant subject, but one that is confronting teachers in increasing numbers. We know the agony and inner panic when faced with young persons who have chosen to die at their own hands or who are contemplating the “ultimate solution” to life’s problems. Teachers themselves are not immune to suicidal thoughts and wishes.

The following guide is meant primarily to bring information to the teaching community so we can get to grips with this pervasive problem and to ease ourselves into the role of suicide prevention through increasing awareness and, as importantly, through the development of a skill called “Active Listening”.



## The role of education


Reluctance to speak openly about suicide and the resulting taboos and myths about suicide predate the modern period.

Historical reactions to suicide have changed in accordance with religious, social, and legal changes in values. Suicide at various times has been considered socially acceptable, a moral evil, or a criminal act. Various societies have seen fit to mutilate the body of a suicide victim, ostracise surviving family members, or cast a shroud of eternal damnation on the victim.

Because of the historically negative reactions to suicide and its victims, guilt, emotional anguish, social embarrassment, self-blame, fear, and superstition still colour reactions to and about suicide.

As teachers, it is imperative that we challenge these historical taboos and myths and replace them with a learned approach to, and compassionate understanding of, those who suffer to the point of taking their own life.

Being human does not require specialised training. All we need is a compassionate, caring attitude, a non-judgmental understanding, and a willingness to listen, mixed with a basic knowledge of the causes, warning signs, and suicide awareness and prevention techniques.



# AN INTRODUCTION TO YOUTH SELF-HARM



Suicide and self-harm in the youth of Ireland is receiving an increasing amount of coverage over the last number of years, and with good reason. Unfortunately, Ireland is one of the most severely affected countries in the EU in this regard. While there is some recent cause for very cautious optimism, there is much work yet to be done.

While depression and self-harm is far more common in females than males, completed suicide remains far more common in males. In the past, one of the explanations given was that while females were more open about describing their feelings, males tended to “bottle them up”, and often resorted to drugs or alcohol to deal with their emotional distress. However, recent work done by the HSE has shown that drug and alcohol misuse is at least as prevalent in teenage females as males, suggesting that we have to look elsewhere for reasons.

While the suicide rate is a very important indicator, by no means does it tell the whole story. The rates of depression in young people are extremely important, because the level of impairment which depression causes can be marked, preventing young people from reaching their potential, and affecting them emotionally, academically and socially.

The issue of self-harm has certainly generated much discussion over recent years, with some research suggesting that children as young as seven years old can engage in such activity. While this is very much the exception, it nonetheless backs up the impression that self-harm in young people is happening at a younger age, and appears to be more widespread. There are many reasons why people engage in self-harm, and it would be wrong to assume that everybody who engages in self-harm is suicidal, and equally it would be incorrect to assume that everybody who engages in self-harm is depressed. Nonetheless, it does represent a worrying act because of its associations.

The most common forms of serious self-harm are cutting (particularly to the arms, but less frequently to the legs, abdomen and torso) or poisoning (particularly with over-the-counter medications), and choking/hanging (which is more common in completed suicides).

## The Four Steps to Help Programme

The Four Steps to Help Programme for Schools has been developed and approved by our Advisory Panel which includes a consultant child and adolescent psychiatrist and has been deemed to be age appropriate for ages 16+.

These school visits are fully funded by the YSPI National School Visits Programme as part of our outreach work and there is no cost to the school or organisation to book and host a talk.

The talks we present are based on our own “Four Steps to Help Programme” which is designed to increase awareness of suicide prevention and to provide information on the support and resources available to anyone who is concerned about suicide or is concerned about a friend or family member.

The Four Steps to Help Programme for Schools focuses on providing simple, factual guidance for suicide awareness and prevention.

The programme emphasises:

- ✦ the importance of awareness of sudden changes in behaviour in friends, family or acquaintances;
- ✦ understanding how to recognise the suicide warning signs, as well as signs of other mental health issues;
- ✦ promotion of 'active listening', giving simple listening skills and exercises which can be used in everyday situations;
- ✦ emphasis on making appropriate family members or other adults aware of concerns so that action can be taken;
- ✦ practical information on dealing with a crisis situation and how to respond.



We all need to know and understand that there is always someone that we can turn to in our life and talk about the issues facing us without fear of rejection if we are prepared to take the first step. What is much more challenging is being prepared to take that first step for someone else; to face possible rejection or ridicule by being prepared to use the Four Step Programme to potentially save a life.

The Four Steps to Help Programme is a simple introduction to suicide prevention and mental health awareness information. **The talks focus on four steps:**



### Step 1: Watching

This step focusses on raising awareness of the suicide warning signs and encouraging participants to watch out for sudden behaviour changes in their friends and family members.



### Step 2: Showing

This step focusses on showing friends and family members that they can discuss any concerns with their friend or sibling, and they can turn to parents and trusted adults for support and advice. This step introduces the Active Listening skillset and provides techniques and examples.



### Step 3: Asking

This step focusses on simple techniques for asking someone how they are feeling and encouraging them to feel safe to talk about their concerns. This step builds on the techniques introduced in step 2.



### Step 4: Helping

This step provides guidance on how to help someone you are concerned about and focusses on getting assistance from a trusted adult or directly from a medical professional. This step also introduces a crisis scenario and provides information on how to deal with the need to get immediate assistance. This step links to the YSPI FreeText Crisis Information service detailed on the back page of this leaflet.

## Why awareness and vigilance are so important

Young males in Ireland between ages 10-24 are more likely to die by suicide than by any other cause based on CSO Statistical Data from 2014. Suicides in 2014 were registered at 10 per 100,000 of the population, an increase on previous years with males accounting for 80% of all registered suicide death (CSO Statistical Data 2014) leading to Ireland having the second highest youth suicide rates in the European Union (Eurostat).



We can bring those rates of youth suicide under control if we reach young people at their most vulnerable ages, if we can spot the early signs of distress and do our best to reduce those stressors as much as possible. As teachers we can greatly assist in that process by being aware, knowing what to look for and encouraging our students to do the same.

## Fighting stigma



One of the most positive influences we can have is by fighting stigma. There is still a huge amount of stigma attached to suicide, and sometimes this can be institutionalised in schools.

Some students have given us feedback that when they wanted to visit a friend who had attempted suicide they were actively discouraged by some teachers from doing so; although the same teachers had been encouraging their students to visit a young student from their school who had been diagnosed with leukaemia. The students couldn't see how the situations differed but they were being given different advice.

None of us are perfect, and we all have prejudices, but it is so important that we allow our students to develop their own attitudes to issues which challenge them, their friends and peers. This is particularly the case with suicide, where we often still see the young person who has lost their way and tried to end their suffering treated as a perpetrator rather than a victim.

## FREQUENTLY ASKED QUESTIONS

### Why do people die by suicide?

People who die by suicide are often having intense feelings of helplessness and hopelessness and may not see any other way out of their emotional pain. It is important to remember that most people who attempt suicide do not really want to die. They simply want to end the pain they are experiencing.

1

### 2 Is it true that people attempt suicide as a cry for help?

The suicide attempt is quite often a conscious or unconscious method for getting others to recognise just how badly the individual is feeling. Yes, suicide attempts are very often cries for help.

### If someone in a family has completed suicide, are other members of the family tempted to try suicide when they have problems?

If someone in a family has completed suicide, other family members may be tempted because suicidal behaviour has been "modelled" for them. However, suicide behaviours are not inherited in families.

3

### 4 Do people ever attempt suicide to "get attention" or to get others to feel sorry for them?

Anyone who attempts suicide in order to get attention desperately needs it. It is tragic when someone feels they need to bargain with their life in order to have their problems taken seriously. Any suicide attempt needs to be taken seriously.

### If a person attempts suicide and fails, what is the likelihood of them trying again?

One of the important warning signs for suicide is a prior attempt. Anyone who attempts suicide once is more likely to try suicide again than those who have never attempted. However, many people who receive licensed professional medical and behavioural health care following a suicide attempt may never become suicidal again.

5



6

### Is it true that people who attempt to kill themselves really don't want to die?

Many people who attempt suicide are ambivalent about life. They want to live and die at the same time. But, as noted in number 1, it is not that the person really wants to die, but rather that death may seem like the only way to end the emotional pain the suicidal person may be feeling. It is the pain they want to end usually, not the life.

### Will a person who is deeply depressed always become suicidal?

While it is true that suicidal feelings often develop in a person who is deeply depressed, the fact that one is depressed does not mean that a person will become suicidal.

7

8

### Does anyone ever impulsively try suicide and then become sorry for making such an attempt?

A person at a particular moment may find the emotional pain being experienced absolutely intolerable. At a given moment, a suicide attempt might impulsively be made which, in retrospect, might be regretted.

### Does taking drugs or alcohol increase one's chances of becoming suicidal?

Taking drugs or alcohol in excess can exaggerate painful feelings to a point where the feelings become intolerable. In such a state, a person might attempt suicide who otherwise would not go that far.

9

10

### Is a person who attempts suicide mentally ill?

All suicidal individuals are not necessarily mentally ill, though many people who attempt or complete suicide may have symptoms of mental illness, the most common being some form of depression. It is important to note that most depression is of a temporary nature and is treatable.

### Is it true that gay teenagers are at higher risk for suicide than teenagers in general?

Studies in the US indicate that gay, lesbian and bisexual youth account for some 30% of all youth suicides, yet constitute only about 10% of the total youth population. Thus, it is clear that such youth are at much higher risk for suicide than the youth population as a whole.

11

12

### How can one help a person who is suicidal?

A person who feels that life is too painful is often feeling very worthless, perhaps unloved, perhaps isolated. Showing such individuals some real caring, by listening to them, accepting their feelings without judgment, by staying close, and getting others to be supportive, can really help. Giving time and really listening to someone in crisis is critical. It may be important to refer the person to a professional medical or mental health worker at some point.

### How does talking about suicide help to prevent it?

Talking about suicide diffuses some of the intensity of suicidal feelings. It helps the person get connected to the help that may be needed. It creates a climate of caring and helps to break through the loneliness and isolation a person may be experiencing. By asking someone in crisis if they are suicidal, we give that person permission to talk about possible suicidal feelings, about which they may otherwise feel they cannot, or should not, talk about.

13

14

### Is suicide or attempted suicide against the law?

At one time suicide or attempted suicide was against the law. In some countries it has only been within the last 20 years that suicide has ceased to be a crime.

### What effects does a suicide have on the individual's remaining family and friends?

15

The survivors of a suicide are left with complex and often confusing feelings of rage, guilt, despair, grief, loss, shame, etc. Recovery from the loss of a loved one by suicide is a very difficult form of grief to resolve, and may never be completely resolved. It has been estimated that every suicide, on average, has a direct, profound emotional impact on 8 to 12 other people. With some 30,000 suicides each year in the EU, there are consequently a huge number of emotionally impacted "suicide survivors".

### Why do some people keep secret the fact of a suicide in the family?

Some people keep the fact of suicide in the family a secret out of fear of being blamed or socially ostracized. Fortunately today, much of the historical stigma of suicide is lifting and people are dealing with suicidal death more directly and honestly.

16

### What are the most common methods used by teenagers to attempt or complete suicides?

17

Lethal methods for attempting suicide by teenagers include guns, hanging, carbon monoxide, jumping, and drug overdoses. Auto accidents account for many deaths, but it is often difficult to determine whether the death is suicide or an accident.

### Do more men or women make attempts on their lives?

Although about three times as many women attempt suicide than do men, about four times as many men complete suicide than do women. This is due to the fact that men use more lethal methods, such as guns or hanging, while women are more likely to attempt suicide by using pills.

18

### Does everybody think about suicide at least once in their lifetime?

19

At some point in their lives, most people have at least fleeting thoughts of suicide, especially in times of personal crisis but it does not mean a person will die by suicide.

## Myths about suicide

### 1. People who talk about suicide seldom mean it and can, in fact, be regarded as low risk to attempt suicide.

Quite often people who talk about suicide do attempt suicide. Most of those who kill themselves have left definite warning signs. One of the major clues is talking about suicide. Many times people make statements about suicidal intentions to see how others respond, to see if anyone cares enough to ask about what is going on. It is important to treat all talk about suicide seriously. The talk may be a cry for help.

FALSE

### 2. The fact that someone has attempted suicide once greatly reduces the risk of a second attempt

A suicide attempt is considered a cry for help. Once an attempt is made, the person is at higher risk for making another attempt or completing suicide. Statistics show that 10 percent of those who attempt suicide will eventually try to kill themselves again, two percent within a year. Many others will make further attempts but survive. Young people are, in many instances, likely to make successive attempts. The elderly, on the other hand, are more likely to die with a first attempt. Of course, not everyone who attempts suicide once will try again.

FALSE

### 3. Women attempt suicide more often than men.

It is estimated that women make 3 times as many suicide attempts as men. However, it is also estimated that men actually complete suicide four times as often as women. In other words, suicide attempters tend to be female, while suicide completers tend to be male. One of the reasons for this statistical disparity is that men tend to use more lethal means, such as weapons, jumping or hanging, than women use when attempting suicide.

TRUE

### 4. Suicide now ranks among the leading causes of youth death in the European Union.

In the European Union during 2015, according to Eurostat, there were approximately 56,000 reported deaths by suicide making it one of the leading causes of death. Males accounted for 43,000 of those deaths or 76%.

TRUE

### 5. Most suicides occur "out of the blue" without any warning signs.

Although suicide often comes as a shock and seems to have happened with great suddenness and without warning, the truth is that there are clear and identifiable warning signs in the majority of cases. The problem is that: a) not all the warning signs are obvious (some are subtle) and b) not everyone knows what the warning signs are. That is why suicide awareness education is so important, so people can both recognize the warning signs and respond to them by helping a person in crisis.

FALSE

### 6. Asking someone if they are thinking about suicide will put the idea in their head.

Asking someone if they are thinking about killing themselves can be a scary thing. However, it is important to know that asking the question will not put the idea in someone's head. It will not reinforce the idea, if the person is already thinking about suicide. It will not cause the person to kill themselves. Many times, after being asked, a person finds it a real relief to talk about it with someone else, someone who cares enough to ask. Asking the question gives the person permission to talk about it. It may well be the first step in getting the person some needed help.

FALSE

### 7. Teenagers rarely attempt suicide, although they may frequently think about it.

In all likelihood, teenagers make more suicide attempts than any other age group. So not only do teenagers frequently think about suicide; they also frequently attempt it (at a rate of about 500,000 attempts per year in the United States). Perhaps two percent of all high school teens have made at least one suicide attempt. One study suggested that half of all teenagers have "seriously considered" suicide by the time they graduate high school. Suicide is a very serious issue for this age group.

FALSE

### 8. Suicide attempts mean a person has ambivalent feelings toward life and death.

Studies indicate that the vast majority (perhaps as many as 95%) of people who attempt suicide either do not want to die or are not sure. They are ambivalent. Part of them wants to live. Part of them wants the emotional pain they are experiencing to stop. While struggling with, often times unbearable emotional pain, they think about or plan a way out, often giving signals of their distress to significant persons around them. Suicidal people almost never want to die; they just want the pain to go away.

TRUE

### 9. Improvement following a suicidal crisis means that the suicidal risk is over.

A large number of suicides occur within the first several months following signs of "improvement" in the condition of a person in crisis. This may well be the time when the person finally has the energy to put morbid thoughts and feelings into effect and possibly attempt suicide. Relatives, friends and health care professionals need to remain especially vigilant during this apparent "improvement" period.

FALSE

**10. Suicide is inherited or runs in the family.**

Suicide is an individual problem but risk factors can include a family history of suicide and other health and behavioural health issues that may or may not put a person at risk for suicide. Other individual and family lifestyle issues can also be risk factors for suicide including the use of drugs and alcohol.

**FALSE**

**11. All suicidal individuals are mentally ill and suicide always is the act of a psychotic person.**

All suicidal individuals are not necessarily mentally ill, nor is suicide always the act of a psychotic person. Many people who attempt or complete suicide may have symptoms of mental illness, the most common symptom being some form of depression. (Most depression is treatable and temporary in nature). Some studies do indicate that between 40 and 60 percent of people who died by suicide were at the time experiencing an episode of depression.

**FALSE**

**12. The incidence of suicide among the poor and deprived is substantially higher than among the advantaged.**

Suicide is quite democratic. It is neither the poor man's curse nor the rich man's disease, but is represented proportionately among all levels of society based on advantage or disadvantage.

**FALSE**

**13. At least half of all people who complete suicide leave notes explaining their action.**

Studies have showed that only approximately 15 to 20 percent of those who complete suicide leave any type of note, and not all notes provide explanations of the person's actions. One of the difficult realities many survivors of suicide (family members and significant others left behind) must come to terms with in many cases, is the fact that they will never really know why someone completed suicide.

**FALSE**

**14. The elderly have the highest rate of suicide for any age group.**

For many people, their so called golden years are anything but golden. Age inevitably brings with it certain losses. Friends and relatives die, careers come to a close, and physical health begins to fail. It can be a difficult and even depressing time of life - a time when many are at heightened risk for suicide.

**TRUE**





# WHAT SHOULD I BE WATCHING FOR?

Teachers are in the unique position to observe students individually or in groups, to sense the emotions seething under the surface, and to monitor subtle changes in behaviour, and to realise that the student in reality might be a hurting human being.

It is possible for teachers, through no fault of their own, to misinterpret a warning sign of suicide or to not even be aware of it. This can be said of the most caring of persons. We were not born with intuitive knowledge. Along with everyone else, we must be trained to discern the messages and signals that our students are sending out.

Someone might ask, "Won't I appear foolish if I misread the signal and get involved in a non-suicidal situation?" That is possible; but if we need to appear foolish to possibly save a life, then it is better than to suffer self-recrimination for doing nothing. The risk of a student dying is more serious than the possible risk of being embarrassed for asking the "wrong" question. If a teacher does raise the question of suicide needlessly, the only reaction possible is a positive one - "This teacher cares."

The lesson that follows is designed to help teachers as well as students become aware of the hidden messages that are silently coming our way from someone in crisis who may well be planning on ending his or her life.



## Some of the underlying causes of youth suicide and self-harm

Much work has been done to try to clarify the reasons why young people kill themselves. We know that young people who are depressed are more likely to go on to complete suicide, but it is important to state that it is still only a relatively small percentage. It is also the case that drug and alcohol misuse is certainly more common in those who die by suicide.

Drugs and alcohol tend to have a two-fold effect. The first is that they, in time, act as a depressant, making low mood more likely. The second is that they tend to decrease inhibitions, and therefore remove the internal controls whereby one prevents oneself from engaging in self-harm or suicide. It is also important to be aware that drug and alcohol misuse can be a marker, in any young person, for increasing levels of unhappiness or distress. It is not unusual for those who are unhappy to turn to drugs or alcohol to give themselves a lift, but, as has been mentioned, the effect is short lived and is replaced by a further deterioration in mood.



When young people self-harm, they give a variety of reasons, but certain themes quickly emerge. The most common of these is the break-up of a relationship, or other peer difficulties.

Another common situation is difficulties at home within the family, and such difficulties can either be of longstanding duration or acute. Less frequently young people describe a build-up of pressure, either academic or otherwise, which generally leads to mounting levels of tension and stress and a feeling that a young person cannot cope. If this happens in the context of a personality which is somewhat perfectionistic and rigid in nature, where a young person does not allow themselves the option of "failure", the result can be catastrophic because a young person's problem-solving competence in such situations becomes significantly impaired, as does their list of possible remedies.

This issue of problem-solving difficulties is a recurrent one in young people because in many cases of self-harm a young person's ability to generate an alternative solution is defective. This happens for a variety of reasons.



The first of these is that, for many young people, their ability to put words to their feelings is at a relatively undeveloped stage. Even though we live in an age where psychological terminology is far more part of the common usage than it was in the past, there is still often a gap between a young person's awareness of the meaning of a term and their ability to apply it to their own situation.

Young people discover, as this ability improves, that they can develop some sense of control by being able to articulate their own internal world, and their developing capacity for abstract thought allows them to discuss these topics in a different way. It is also important to remember that our brains work in a way that makes it more difficult for us to think creatively and flexibly about a difficult situation if we are emotionally aroused. Our ability to generate solutions, to estimate risk, to predict the responses of others, and to manage ourselves through such difficult situations is never as good when we are agitated or distressed as when we are calm. We tend to "catastrophise" whereby we imagine the worst possible outcome and therefore react, which can involve self-harm.

Self-harm also develops as part of an on-going pattern. While it may start in a situation where people feel acutely distressed, it often becomes a means of regulating one's levels of discomfort. Hard though it is to believe, young people who cut themselves while highly distressed describe, not a sense of pain, but a sense of relief. There are many theories as to why this may happen, but fundamental to understanding this dynamic is to realise that, for a young person, cutting themselves may not always be a painful or a distressing act.

Essentially, young people harm themselves (either by cutting or by overdosing, or by drug or alcohol misuse) to get rid of unpleasant feelings. It can create a cycle whereby they believe that the only way to get rid of such feelings is to carry out that specific act, and a pattern ensues. That is the reason why many people who harm themselves on one occasion go on to repeat such an act. It is by no means always a suicidal gesture, and while a very unhelpful way of coping, must be seen in context. In this regard, cutting is the most common method of self-harm which becomes repetitive.

Overdoses tend to be more serious, and are far more likely to have lethal intent. Hanging and choking are almost always lethal with respect to intent and must be taken very seriously. The one exception to this is that many young children, either in their early teens or perhaps younger, can engage in either breath-holding or asphyxiation games in order to induce an altered state of consciousness. While this does not, generally, have lethal intent, it is quite likely that, in situations where people carry this out alone (by using a ligature) they may be unable to release the ligature in time, and this has led to loss of consciousness, and even death.

## Cyber Bullying

One of the reasons frequently given by young persons who harm themselves is social exclusion by peers or indeed overt bullying. Traditionally, this took the form of either physical aggression or a form of relational victimisation whereby a young person was in some way excluded or teased. With the dramatic increase in communication technologies, the means by which this can be perpetrated have increased, in a way which is extremely difficult to monitor.

Under the umbrella term of cyber bullying come a variety of forms of victimisation which include the use of mobile telephones and computers.





These include abusive messages, spreading rumours, posting photographs either via mobile phone or on social network sites. A further and sinister development is the filming of physical aggression perpetrated on the young person and its transmission to others.

It is extremely difficult to police these forms of communication, in many ways because the technological awareness of teenagers is often far ahead of that of their parents.

In addition, while previously there was just one telephone in a house, and nobody would dare ring after 9.00 p.m. or so, young people with mobile phones or computer access can be busy communicating well into the early hours, when others are asleep. One of the dreadful consequences of being the victim of such bullying is the fact that 'nowhere is safe'. While previously one knew where to avoid, the sense of vulnerability as a consequence of being cyber bullied is much greater.

It is extremely important that young people learn both how to protect themselves and also to report victimisation of others. What can appear to be a "prank" or "teasing" can be highly distressing and has certainly lead to numerous episodes of self-harm, and indeed worse.

## Suicide Clusters

Suicide clusters are a recent phenomenon, and cause tremendous concern. With the mushrooming of reality television has come a level of exhibitionism which was previously not in evidence. It has led to a situation where many vulnerable young people lay bare their inner-most thoughts and emotions for an audience, particularly when distressed.


When one adds to this the possibility of networking with other like-minded individuals on the internet, one can see very clearly how suicide clusters can form. While there are still a number of extremely worrying situations where friends will engage in some sort of suicide pact, the geographical dispersal of those involved in internet-based suicide clusters makes it extremely difficult to penetrate and to prevent.

Again, as with cyber bullying, because of the technological awareness of teenagers compared with most adults, the only way of monitoring such situations is by being able to address these topics in a very open way where young people are not blamed for the sites they frequent, in order that they can feel safe to disclose things that particularly concern them.

It is also true that, albeit rarely, clusters can develop in a single geographic location. It tends to be among a disaffected peer group, and it is often facilitated by drug and alcohol use. It is certainly the case that, with any situation where a young person dies by suicide, the level of suspicion must be high that others may be considering the same thing. In such situations, one finds that groups of youths tend to cluster together, and it feels as if adults are excluded.

Nonetheless, what does often happen is that, while teenagers may not talk to adults about their feelings, they certainly may talk about others within their group, and by networking together, a group of concerned adults (including teachers, parents, youth leaders etc.) can build up quite a detailed picture of the various levels of vulnerability within the group.

Furthermore, in the wake of such bad news, a sense of unresolved loss in others may often be triggered, and therefore the index of suspicion among others, not intimately connected with the core group, must still be high. In such situations, the National Education Psychology Service has a very coherent plan for how best to manage a situation within a school setting, and schools or other Youth organisations are often well advised to seek outside advice on how to manage such situations.



It is a time of intense vulnerability and anxiety, particularly among adults. The tendency for young people to idealise the deceased is very common, and, because one is always so reluctant to speak ill of the dead, it is very difficult to challenge. In time, such intense feelings diminish, but vigilance must remain high for those who continue to show depressive features.

### Mental Health Warning Signs to watch out for:

MAY INCLUDE BUT ARE NOT LIMITED TO:



## WARNING SIGNS AND BEHAVIOURAL CHANGES

While suicide is, by its nature, a terminal act, one must be vigilant for the tell-tale signs of a lowering of mood or of other at-risk behaviours. Unfortunately, not every incident of suicide can be predicted, as, even with the benefit of hindsight, it appears clear that some young people tragically kill themselves in an impulsive manner which could not have been foreseen, even by their nearest and dearest. However, it makes it all the more important that we pay very close attention to signs of depression or increasing impulsivity which can lead to timely interventions where appropriate.

Depression is a condition which previously had been considered unusual in teenagers and vanishingly rare in pre-teens. We now know that the incidence of depression, while uncommon, is very much a concern in pre-teens, and rising through adolescence. It is more common in girls than boys, and its incidence increases with age, until it reaches adult levels.

Depression is characterised by low mood, increasing irritability, social withdrawal, poor concentration, and is often accompanied by alterations in sleep pattern and appetite. The thinking patterns which are common in depression are self-critical, finding fault with many things, pessimistic regarding the future and discounting anything which may appear to be of value or benefit to the person. Essentially, it is the very opposite of "rose tinted glasses".





While most cases of depression are mild to moderate, and resolve within a month or two, some cases become more serious. Teenagers with depression create difficulties for themselves by virtue of their irritability, their deteriorating school performance, their social withdrawal and their resultant behavioural difficulties and lack the flexibility to get themselves out of such situations, because of their low mood. Therefore they can become alienated from family, and sometimes from their friends.

For teenagers, who take so many of their values and their points of reference from their peer group, to fall out with friends, particularly when they have already fallen out with their families, causes huge difficulties, thereby perpetuating and indeed deepening the cycle. In such cases, particularly when teenagers become agitated and angry, self-harm is often an accompanying concern.

One can see, for such teenagers, how drugs or alcohol might offer a temporary reprieve. Nonetheless, it remains that most episodes of drink and drug use by teenagers is of an experimental type, at least initially, and then often part of a peer activity. It is certainly not the intention to make excuses for it, but merely to point out that it is common, and it is usually not associated with mental illness.

Many teenagers, especially boys, are quite impulsive by nature. This can often be seen in the context of young people with ADHD (Attention Deficit Hyperactivity Disorder) but is not restricted to this. For such teenagers, who struggle with deferred gratification, the likelihood of an impulsive act of self-harm is greater than for the general teenage population.

While in some ways such acts are more easily excused, and indeed occasionally explained away because of the impulsivity, when one thinks further about it, one realises that the level of impulsivity actually makes it more difficult to plan prevention strategies compared to those whose acts of self-harm are the result of the more protracted planning.



## Main Indicators

Probably the most important indicator that a student is struggling with something will be a sudden change in their behaviour. So, for example, the student who is very outgoing in class, the class clown, suddenly became very quiet, very withdrawn, you might be grateful but you would probably think 'That's strange. I wonder what's up there'.

If you are involved with teams then there is usually somebody in the team who is first at practice, helps with the equipment, very keen. If they were suddenly to turn around and say that were not going to practice, that they were "not interested", "can't be bothered", you would probably take notice of that change anyways.

- ✦ The student who has always been very keen on studying and suddenly they're cutting school, they just don't care.
- ✦ The student who was always very interested in looking after their body, who has always been eating salads and suddenly, they've been down to the chipper every lunch time stuffing their faces.
- ✦ The student in the class who has always been academic, very keen to be top of the class. Suddenly they're always late or missing school....why?



It's these kinds of sudden changes in behaviour that you will already take note of, but you might not have been aware of the wider potential issues that this behavioural change may indicate.

## General Warning Signs

If someone is seriously depressed and thinking of attempting suicide there are often warning signs that family and friends can pick up on. Noticing and acting upon these warning signs could save a life. Most people who are considering suicide are willing to talk about their problems if someone shows they care. Don't be afraid of discussing the subject with someone you think may be suicidal. Talking about suicide won't 'plant the idea' in someone's head. This is a myth. If you are wrong, you're at least showing a friend you care. If you are right, you could save their life.

Sometimes stress or a traumatic event like bereavement can trigger suicidal thoughts in a vulnerable person. For this reason it's important to ask a friend who is going through a tough time how they are coping and if they need some support. Having someone to talk with can make all the difference.



### Warning signs can include but are not limited to:

- ✦ Withdrawing from family and friends.
- ✦ Having difficulty concentrating and thinking clearly.
- ✦ Sleeping too much or too little.
- ✦ Feeling tired most of the time.
- ✦ Gaining or losing a significant amount of weight.
- ✦ Talking about feeling hopeless or guilty.
- ✦ Talking about suicide or death.
- ✦ Self-destructive behaviour like drinking too much or abusing drugs.
- ✦ Losing interest in favourite things or activities.
- ✦ Giving away prized possessions.
- ✦ Mood swings.

### Additional warning signs that a teen may be considering suicide:

- ✦ Change in eating and sleeping habits.
- ✦ Withdrawal from friends, family, and regular activities.
- ✦ Violent or rebellious behaviour, running away.
- ✦ Drug and alcohol use.
- ✦ Unusual neglect of personal appearance.
- ✦ Persistent boredom, difficulty concentrating, or a decline in the quality of schoolwork.
- ✦ Frequent complaints about physical symptoms, often related to emotions, such as stomach-aches, headaches, fatigue, etc.
- ✦ Not tolerating praise or rewards.

## IMPORTANT NOTE

**If someone mentions suicide, take it seriously. If they have expressed an immediate plan, or have access to prescription medication or other potentially deadly means, do not leave them alone. Get help immediately.**

## Teen specific warning signs

Talking about suicide	Any talk about suicide, dying, or self-harm, such as <i>"I wish I hadn't been born," "If I see you again..."</i> and <i>"I'd be better off dead."</i>
Seeking out lethal means	Seeking access to guns, pills, knives, or other objects that could be used in a suicide attempt.
Preoccupation with death	Unusual focus on death, dying, or violence. Writing poems or stories about death.
No hope for the future	Feelings of helplessness, hopelessness, and being trapped ( <i>"There's no way out"</i> ). Belief that things will never get better or change.
Self-loathing, self-hatred	Feelings of worthlessness, guilt, shame, and self-hatred. Feeling like a burden ( <i>"Everyone would be better off without me"</i> ).
Getting affairs in order	Making out a will. Giving away prized possessions. Making arrangements for family members.
Saying goodbye	Unusual or unexpected visits or calls to family and friends. Saying goodbye to people as if they won't be seen again.
Withdrawing from others	Withdrawing from friends and family. Increasing social isolation. Desire to be left alone.
Self-destructive behaviour	Increased alcohol or drug use, reckless driving, unsafe sex. Taking unnecessary risks as if they have a "death wish."
Sudden sense of calm	A sudden sense of calm and happiness after being extremely depressed can mean that the person has made a decision to die by suicide.

## General risk factors

### LOSSES:

#### 1. The break-up of a romantic relationship

For an adolescent the loss of such a relationship is traumatic in many cases. His or her world has come crashing down. Behind many a macho exterior or sour grapes attitude is a sensitive and hurting young person. Trite expressions like "Things will get better in time" or "There are other fish in the sea" show no sensitivity for the hurt the young person is feeling and deny that the pain is real.



#### 2. The death of a loved one

The pain of separation by death can be so great that the young person might be driven to join that person in death. Furthermore, the grief process often does not include the young person in the family. Many adults do not consider the possibility that the grief that a young person is experiencing at the death of a close family member is as profound as their own.



#### 3. The death of a pet

Consider the teenager whose only true listener is the dog. The dog is there to listen and to love and to never pass judgment. And if that dog should die?



#### 4. The loss of a job

For many teenagers, "job" means maturity and independence. Take away the job? What happens to the independence?



## 5. Losing face

Consider the boy who publicly stated he was aiming to be a team captain and didn't make it. Consider the teen who wanted to attend a prestigious college but got a rejection instead, and everyone knows it.



## 6. Divorce

The loss of a parent through divorce is more traumatic than is commonly admitted. Many teenagers feel responsible for the break-up of the marriage. The imagined or actual fear of a possible divorce is also tremendously painful for the teenager.



### PRESSURES:

#### 1. School Pressure

The need to achieve high marks, time to accomplish several major assignments simultaneously, involvement in too many extracurricular activities, demands of school sports, college applications.

#### 2. Peer Pressure

The need to find acceptance, group morals, conformity to clothing styles, drugs, alcohol, sex, and bullying to name just few.

#### 3. Parental Pressure

Success, money, the right college, the right friends, good marks, conflict between the need to control and the need to be independent, marital problems between parents, "get a job", clothing, music, the parent who wants to be a "friend", lectures rather than examples.

### LOW SELF-ESTEEM:

#### 1. Physical Unattractiveness

Consider the young man who thinks that physically he does not match his peers. Consider the young lady who thinks she's plain and homely. Consider the effect of skin blemishes at debs' time.

#### 2. Never the first

Consider the young man or the young lady who always feels like a second choice when it comes to dating or being chosen for anything.

#### 3. Sexuality

Consider the pain and agony of the teenager who is caught between the two worlds of sexuality and who is terrified to speak to anyone about this for fear of ridicule. Consider the young person whose fear of being homosexual is based on a lack of fundamental sexual knowledge.

#### 4. Clothing

Consider the teenager who, influenced by the media blitz and by teenage styles, judges importance or lack thereof by the type of clothes he or she is forced to wear.

#### 5. Physical Disability

Consider the teenager who must not only cope with a physical problem, but also with the unkind remarks and glances of others.

#### 6. Academic Disability

Consider the teenager whose older brother or sister was a "genius" and is constantly reminded of the difference between them.



# LACK OF COMMUNICATION AND LACK OF HOPE:

## 1. Isolation and loneliness

Many teenagers feel so isolated and alone that they are convinced that there is no one to help them and that no one really cares. Whether this is true or not is irrelevant. What matters is that this is how they perceive it, and so they suffer in silent isolation.



## 2. Without a future and hopelessness

Consider the teenager who instead of looking to the future with expectation is overwhelmed with a sense of hopelessness. All hope in the future has been lost.



## Teen specific risk factors

The teenage years can be emotionally turbulent and stressful for all teenagers. Teenagers face pressures to succeed and fit in. They may struggle with self-esteem issues, self-doubt, and feelings of alienation. For some, this leads to suicide. Depression is also a major risk factor for teen suicide.

Other risk factors for teenage suicide include:

- Childhood abuse
- Recent traumatic event
- Lack of a support network
- Availability of means of suicide
- Hostile social or school environment
- Exposure to other teen suicides



## The Dangerous Calm

The “Dangerous Calm” is a recently highlighted mental health phenomenon. A student may show sudden changes of behaviour, often to quite a significant extent. The student then reverts back to their normal disposition and behaviours.

In 95% of cases this would be perfectly normal as everyone has times of depression, sadness, ill-temper etc. But in 5% of cases this reversion to normal behaviour can mask a life decision by the student. Some students can make a decision in their depressed state that appears to provide a solution to all their problems and, as a result of that decision, their behaviour normalises.

Unfortunately that decision can often be a decision to take their own life, and because that decision becomes internalised and part of a coping strategy, it can remain dormant unless challenged or diagnosed. This is why it is so important that sudden changes in student behaviour are noticed, reported and acted upon.



# BASIC GUIDELINES ON HOW TO HELP

When an adult becomes aware of a young person's emotional distress or their self-harm, it comes about either because an adult has, through vigilance or information received, developed an opinion that a young person is at risk, or else a young person has taken the initiative in discussing their concerns with an adult.

Firstly, it is worth considering what the qualities are in an adult that makes a young person feel they can confide in them.



## Tolerant and accepting

The issue of stigma regarding psychological difficulties is one which has been widely discussed, and is very much seen as a barrier to intervention. Young people are far more likely to approach those whom they know have a tolerant and accepting attitude towards psychological difficulties rather than those whom they perceive to be disparaging or intolerant of such problems. Therefore, the way in which one discusses or talks about psychological problems already sets the scene for whether or not a teenager is likely to approach in case of difficulties.

## Respect

Another significant point is the central importance of the respect which the adult shows to the young person. It cannot be over emphasised that for a young person to describe their own internal world at a time when they feel close to crisis takes a huge amount of courage, and the response of the adult is crucial.

## Not judging or trivialising

The most important aspect is to listen carefully and calmly without judging and without jumping to conclusions. Not only must the adult respect the young person, but they must also respect their account of their difficulties. In other words, one cannot trivialise the symptoms, cannot say, in a dismissive way, that everybody experiences such symptoms, or indeed that the young person has much to be thankful for and aren't there those who have much greater difficulties.

## Allow the young person to talk

It is not a time, when listening to a young person's story, for drawing premature conclusions or cutting them short. The young person is the expert in their own story. It is also worth remembering that young people, in describing such symptoms, may not always feel ready to describe their most pressing concerns, and often, especially if there are particular on-going stressors, may provide a "test case" of such difficulties in order to see how the adult responds.

## Don't guarantee confidentiality

In situations where young people describe episodes of abuse, either physical sexual or emotional, it is absolutely critical that the adult does not give a guarantee of confidentiality to the young person, however much the young person seeks it. To do so, while often based on compassionate grounds, serves to compromise the adult, to undermine the rights of the parent, and indeed on occasion to thwart due process (when the issue is more appropriately dealt with by the civil authorities).

## Listening

In a situation where an adult forms the impression, either from their own observations or from information passed on from others, that a young person is particularly low in mood, or at risk of self-harm or indeed suicide, an approach needs to be made to the young person.

It may well be that the adult who has formed the impression is the best person to do so, or perhaps there is another adult with a more developed relationship with the young person who might be more appropriate. In either event, once the initial approach is made, the importance of listening empathically and without judgement to the young person's answers is critical.



## Discussion does not cause suicide

There is a diminishing, although unfortunately still prevalent, mistaken belief that discussing the topic of suicide is merely implanting this into the young person's mind. This is not true.

It is important that, in situations where it appears that it needs to be asked, the adult does not shy away from asking very specifically about whether or not the young person feels their life is no longer worth living, whether or not they have in any way harmed themselves in the past, or whether they have plans to do so in the future. It is crucial that the adult can tolerate the young person's distress, because this is very containing and comforting for the young person.

## Under 18s

In either event, if one is dealing with a young person under the age of eighteen, it is essential that the young person's parents be first informed. The one exception to this is if there are sufficient grounds for concern that a young person has been the victim of some form of abuse at home and that his or her parents may not have the young person's best interest at heart, then one should approach the civil authorities, most often the HSE Child Protection Team.

## Getting support

However, in the more usual situation, it is important that young people can be reassured, in so far as possible, that their feelings are both validated but also that help is available, and that advice can be given on whom to contact.

In situations where there is a level of unhappiness without any concern regarding self-harm or suicide, then there may well be people locally, either within schools, youth clubs or other organisations, who can support the young person. If there are concerns regarding self-harm or significantly lower mood, then parents are best advised to discuss matters with their family G.P. who will know both the family history and also the range of appropriate local options.



## Helping students – DO'S

- 👍 Do befriend when appropriate
- 👍 Do consider the possibility of suicide
- 👍 Do focus on the pain
- 👍 Do ask if suicide is on their minds
- 👍 Do get involved
- 👍 Do allow them to express feeling
- 👍 **DO LISTEN**
- 👍 Do make life an option for them
- 👍 Do be non-judgmental
- 👍 Do get help and support for yourself as well
- 👍 Do stay with the person at risk

## Helping students – DON'TS

- 🗨️ Don't lecture or moralise or give advice
- 🗨️ Don't think it is a passing phase
- 🗨️ Don't brush off feelings with inane remarks
- 🗨️ Don't be afraid that you will instil the idea of suicide
- 🗨️ Don't do nothing
- 🗨️ Don't trample on feelings
- 🗨️ **DON'T TALK TOO MUCH**
- 🗨️ Don't dare them to follow through with suicide
- 🗨️ Don't react verbally or physically with shock
- 🗨️ Don't go it alone
- 🗨️ Don't leave the person alone
- 🗨️ Don't promise not to tell anyone

## Mental Health Awareness Programmes



Some schools are now finding a positive benefit in incorporating suicide awareness and prevention into a wider mental health programme for the school. Mental Health weeks are a good way of doing this.

Other areas of mental health can then be added such as nutrition, exercise, online behaviour, stress management etc. This makes the “heavier” subjects such as depression, mental illness, addiction and suicide more palatable for the students.

In the case of suicide awareness we can only provide our programme to students aged 16 upwards so generally transition, 5th and 6th years. This would be a good guideline age according to our medical panellists.

## Availability of staff

Feedback from students often mentions that they would like to talk with a teacher but “they’re always busy”, “someone is always with them” or “they’re not very approachable.”

In schools with guidance or pastoral services they will already usually provide an open-door policy. Other schools have found that having a teacher available at certain times of the day in a quiet office or small classroom can provide students with an opportunity to discuss problems without their friends or classmates knowing about it.

## Providing students with coping skills



Perhaps the most important subject we can teach our students is how to cope with the challenges that life presents to each and every one of us. Techniques for dealing with stress and emotions are learned through the modelling of behaviour, trial and error, practice, and more practice through life experiences, and the process of maturing.

Students must be made to recognise that they need to constantly develop techniques and strategies for coping, and to continue to build the necessary skills throughout their lives. This takes a conscious effort, careful thought and practice. Each individual develops his or her own methods and styles of coping with difficulties and stress in their lives. They are as personal as each individual's personality. As we guide and support our students' educational growth and development, so too must we guide, support and reinforce the development of good positive coping skills in our students. While we cannot and should not make decisions for our students, we can help them to become aware that they have "choice" and "control" in their lives.

With choice and control come the ability to learn and develop positive ways of coping with the stresses of life, which are inevitable.

Our task then becomes one of facilitating the exploration of healthy, positive coping techniques, providing support, helping students to grow and mature and helping them in their quest to acquire the "tools" to cope with their emotions and life situations.

## What can I do for my students?

- ✦ Be open to discussing difficult or "taboo" subjects with age-appropriate teens
- ✦ Don't be afraid to use the words: suicide, self-harm, cutting etc.
- ✦ Don't be afraid to challenge teens
- ✦ Be aware of the new challenges facing young people
  - Cyber-bullying
  - Online Blackmail
  - Coercion & intimidation
  - Sexualisation by peers
  - **Be approachable**



# THE ROLE OF THE LISTENER

For teachers the role of listener can be a bit unnerving. As teachers we are much more used to talking, to taking charge of a group and being the source of information. In terms of assisting a student who wants to express themselves to us we need to become more passive.

The funny thing is that listening seems to be so challenging when actually it can be quickly and easily learnt by using the techniques of Active Listening. Active Listening is widely used by helplines such as the Samaritans as it allows a consistent approach, established empathy but, importantly, also allows the listener to keep distance between themselves and the person who wants to talk.

## General guidance

It is actually ok to ask students about their mental state. We need to know where their thoughts are going even though it may seem challenging.

So how do you ask a student about their thinking?

- 👍 **DO be yourself.** Let the person know you care, that he/she is not alone. The right words are often unimportant. If you are concerned, your voice and manner will show it.
- 👍 **DO Listen.** Let the person unload despair, ventilate anger. No matter how negative the conversation seems, the fact that it exists is a positive sign.
- 👍 **DO be sympathetic, non-judgmental, patient, calm, accepting.** Your family member is doing the right thing by talking about his/her feelings.
- 👍 **DO offer hope.** Reassure them that help is available and that these dark, possibly suicidal feelings, are temporary. Let the person know that his or her life is important to you.



If the person says things like, "I'm so depressed, I can't go on," ask the question: "Are you having thoughts of suicide?" **You are not putting ideas in their head; you are showing that you are concerned, that you take them seriously, and that it's OK for them to share their pain with you.**



- 👎 **DON'T argue with them.** Avoid saying things like: "You have so much to live for," "Your suicide will hurt your family," or "Look on the bright side."
- 👎 **DON'T act shocked, lecture on the value of life,** or say that thinking these thoughts is wrong.
- 👎 **DON'T promise confidentiality.** Refuse to be sworn to secrecy. A life could be at stake and you may need to speak to a health professional in order to keep your teen safe. If you promise to keep your discussions secret, you may have to break your word.
- 👎 **DON'T offer ways to fix their problems,** or give advice, or make them feel like they have to justify their feelings. It is not about how bad the problem is, but how badly it's hurting your friend or loved one.
- 👎 **DON'T blame yourself.** You can't "fix" someone's depression, only support them, seek professional assistance and offer love.



## Active Listening

"Active Listening" is simply the offering of friendship by one ordinary human being to another at a time of crisis or loneliness. An Active Listener doesn't need to have professional status or authority, but is simply a fellow human being who cares. The purpose of Active Listening is to listen, accept, care and empathise.



### LISTEN

Allowing the person with a problem to express and to talk without being judged.



### ACCEPT

Allowing the person to stay in neutral and accept their feelings as they are.



### CARE

Allowing one human being to reach out to another human being with respect.



### EMPATHISE



Allowing the listener to hear where the speaker is coming from and allows us to be sensitive to another's feelings or ideas **even when we don't agree.**

The purpose of Active Listening is not to give advice, instruct, solve problems, or judge. It is to respect the worth and value of another human being through Listening, Accepting, Caring, and Empathising.

## Characteristics of good active listening

A good Active Listener is someone who:

### DOES

-  listen more than talk
-  direct the conversation to the painful feelings
-  have compassion for sufferer
-  risk being foolish
-  attempt to be available at all times
-  remain willing to share another person's pain
-  respect confidences
-  listen
-  accept
-  empathise

### DOES NOT

-  offer opinion or judgments
-  belittle or minimise concerns
-  discuss one's own problems
-  give advice
-  express shock or surprise
-  patronise or probe
-  offer platitudes and clichés
-  make promises that cannot be kept
-  interpret, lecture or diagnose
-  say "I know just how you feel."
-  fail to pay attention or care



## Attitude of the listener

### YOU ARE LISTENING TO ME WHEN...

**You come quietly** into my private world and let me be me;

**You really try to understand me** even if I'm not making much sense

**You grasp my point of view** even if it goes against your own sincere convictions;

**You realize** the hour I took from you has left you a bit tired and drained;

**You allow me** the dignity of making my own decisions even though you think they may be wrong;

**You do not take my problem** from me, but allow me to deal with it in my own way;

**You hold back your desire** to give me good advice

**You do not offer me religious solace** when you sense I am not ready for it;

**You give me enough room** to discover for myself what is going on;

**You accept my gift** of gratitude by telling me how good it makes you feel to know you have been helpful



### YOU ARE NOT LISTENING TO ME WHEN...

**You do not care** about me;

**You say you understand** before you know me well enough;

**You have an answer** for my problem before I've finished telling you what my problem is;

**You cut me off** before I have finished speaking;

**You find me boring** and don't tell me;

**You feel critical** of my vocabulary, grammar or accent;

**You are dying** to tell me something;

**You tell me about your experience** making mine seem unimportant;

**You are communicating with someone else** in the room;

**You refuse my thanks** by saying you haven't really done anything.

# IF YOU CANNOT HELP A STUDENT

So what happens if a student comes to talk to you and you are just not comfortable with the idea; you might not be in a good place yourself; you might feel it isn't appropriate; or you just might not relate well to the particular student. For some reason we have noticed that students will often come looking for support from teachers who they have the most confrontation with, which may be because they perceive that teacher as strong or they might even be seeking "permission" to talk about something that is bothering them.

If you find yourself in a position where a student wants to talk with you and you cannot help them then there is some guidance available:

1. Bear in mind that the student may have a serious issue to report
2. Don't let the student start telling you the problem and then cut them off.
3. Be upfront that you can't assist and tell them why, gently.
4. Tell the student who can assist them.
5. Take the student to that person; don't just send them off. That can be seen as rejection.
6. The student came to you for a reason; offer to stay with them while they talk to the other teacher or counsellor.
7. Make sure that you report the fact that the student came to see you so that there is a record.
8. Always follow your school's child protection policy.

## POLICY MATTERS

It is vital that child protection and welfare is always at the heart of any programme.

### Student Protection


Teachers should always follow their school's child welfare policies at all times. If anything in this guide contradicts your local policies or you are unsure about how they should be implemented then you should discuss this with your principal.

### Self-protection

Teachers should always follow their school's child welfare policies at all times. If anything in this guide contradicts your local policies or you are unsure about how they should be implemented then you should discuss this with your principal.



# TYPES OF INTERVENTION RESOURCES



## Examples of school intervention resources

Social Workers

Principal, Deputy Principal,

Teachers and other Staff

Peer Listening Group

School Nurse

## Examples of outside of school intervention resources

Trusted Family Member or Friend

Family Doctor

Community Mental Health Resources

The Samaritans or other helpline

Private Therapist

Counsellor

Clergy



# SUMMARY

It is important to remember that as many as one in four of the population at some point will experience symptoms of depression, and that as many as 10% of children and adolescents will experience impairing psychological symptoms during their childhood years, so these things are by no means uncommon. It is also the case that, for the vast majority, a full recovery is achieved.

However, there are those whose symptoms are severe and impairing who may need quite active intervention in order to allow them to return to their normal activities and regain their developmental trajectory. It is also important to remember that, in many situations, mental health difficulties may not be preventable. The most important aspect is how they are dealt with and this is the hallmark of mature organisations, whereas the less evolved organisations, and by this I include families, schools, clubs etc., have a tendency to disregard any such difficulties, assuming that these things could not happen to them, thereby ignoring them and preventing the appropriate interventions where necessary.



## FreeText HELP to 50015

Texting 50015 is free of charge from any network and you can text this number even if you have no credit. This service is fully confidential and YSPI has no information on mobile numbers that use the FreeText service

### For information on:

- **Emergency Contacts**
- **National Helplines**
- **Local Support Groups**
- **Directions to your nearest GP or out of hours clinic**
- **Directions to your nearest safe place or refuge**

 [www.ineedhelp.ie](http://www.ineedhelp.ie)

**Always call 999 if someone is seriously ill or injured, and their life is at risk.**

**Gardaí / Ambulance / Fire call 999 or 112**





# YOUTH MENTAL HEALTH AWARENESS & SUICIDE PREVENTION

GUIDANCE AND INFORMATION FOR TEACHERS AND LECTURERS

**YOUTH SUICIDE PREVENTION IRELAND**

Registered Charity 20070670



 83A New Street  
Killarney, Co Kerry V93 FR59

 1800 828 888

 [admin@yspi.ie](mailto:admin@yspi.ie)

 [schools.yspi.ie](http://schools.yspi.ie)